



## Choosing a method for correcting anxiety in children aged 4–12 years before dental treatment

Irina A. Nikolskaya<sup>1</sup> , Evgenia N. Anisimova<sup>2</sup> , Igor S. Kopetskiy<sup>1</sup> , Natalia Yu. Anisimova<sup>2</sup> , Pavel A. Voronin<sup>1</sup> , Valeria A. Katyukhina<sup>1</sup>

<sup>1</sup> Pirogov Russian National Research Medical University, Moscow, Russian Federation

<sup>2</sup> Russian Biotechnological University, Moscow, Russian Federation

✉ doknikolskaya@gmail.com

### Abstract

**INTRODUCTION.** Many children have anxiety and fear at the very thought of visiting a dentist. Children do not make contact with a doctor, refuse to perform dental manipulations, express their emotions vividly, or withdraw into themselves. In addition, in a state of anxiety, the work of many organ systems changes and an increase in heart rate is the most common indicator. As a result, such patients refuse to visit the dentist, which worsens the state of oral health. Alternatively, dental procedures for unadapted patients are performed under anesthesia or sedation. But doctors do not have a proven algorithm for diagnosing a child's anxiety level and how to correct this condition for safe and effective dental treatment.

**AIM.** Clinical and functional justification of the choice of a method for correcting anxiety in children aged 4–12 years in an outpatient dental appointment

**MATERIALS AND METHODS.** From 2019–2024 years, a study was conducted, in which 986 children aged 4–12 years participated. Before the start of the examination, general somatic and psychological anamnesis of patients was collected. The level of anxiety of children from 4 to 6 years of age was determined using the “test of child anxiety” by R. Temml, M. Dorka, V. Amen; from 7 to 12 years of age—the “Scale of explicit anxiety CMAS” in the adaptation of A.M. Parishioners. Hemodynamic parameters (HR and SpO<sub>2</sub>) were measured using a Choice MMed 300C5 pediatric pulse oximeter. All patients underwent a dental examination of the oral cavity and the degree of caries activity was determined by T.F. Vinogradova. Based on the results of the examination methods, a decision was made on the further strategy of working with the patient (treatment with the support of an anesthesiologist or the use of art-therapy as a method of psycho-emotional correction).

**RESULTS.** According to the results of psychological testing, all patients were divided into three groups: low-level (19.2%), medium-level (52.4%) and high-level anxiety (28.4%). Almost all children with low levels of anxiety had their heart rate within the age norm (86.9%). This group also showed the highest percentage of patients with compensated caries activity (51.9%). Children with a high level of anxiety showed moderate and severe tachycardia (38.8% and 61.2%, respectively), and 62.2% of the subjects in this group showed a decompensated degree of caries activity. Patients with low levels of anxiety required less psychological correction, than children with medium and high levels. Based on the results of our study, a computerized program “Determining the method of correcting anxiety in children before dental treatment” was developed, which will help dentists determine the tactics of working with children's patients before dental manipulations.

**CONCLUSIONS.** Thus, to improve the quality and safety of dental care for children, on the basis of a comprehensive psychological and functional examination, a digital program “Determining the method of correcting anxiety in children before dental treatment” was developed.

**Keywords:** anxiety, adaptation, heart rate, art therapy, degree of caries activity

**Article info:** received – 06.03.2025; revised – 27.04.2025; accepted – 15.05.2025

**Conflict of interest:** The authors report no conflict of interest.

**Acknowledgements:** There are no funding and individual acknowledgments to declare.

**For citation:** Nikolskaya I.A., Anisimova E.N., Kopetskiy I.S., Anisimova N.Yu., Voronin P.A., Katyukhina V.A. Choosing a method for correcting anxiety in children aged 4–12 years before dental treatment. *Endodontics Today*. 2025;23(2):312–317. <https://doi.org/10.36377/ET-0089>

## Выбор метода коррекции тревожности у детей в возрасте 4–12 лет перед стоматологическим лечением

И.А. Никольская<sup>1</sup> , Е.Н. Анисимова<sup>2</sup> , И.С. Копецкий<sup>1</sup> , Н.Ю. Анисимова<sup>2</sup> , П.А. Воронин<sup>1</sup> , В.А. Катюхина<sup>1</sup>

<sup>1</sup> Российский национальный исследовательский медицинский университет имени Н.И. Пирогова, г. Москва, Российская Федерация

<sup>2</sup> Российский биотехнологический университет, г. Москва, Российская Федерация

✉ doknikolskaya@gmail.com

### Резюме

**ВВЕДЕНИЕ.** Большинство детей испытывают тревогу и страх уже при одной мысли о посещении стоматолога. Они избегают контакта с врачом, отказываются от проведения стоматологических манипуляций, бурно выражают свои эмоции либо, напротив, замыкаются в себе. Кроме того, в состоянии

© Nikolskaya I.A., Anisimova E.N., Kopetskiy I.S., Anisimova N.Yu., Voronin P.A., Katyukhina V.A., 2025

тревоги изменяется работа многих систем органов, а учащение сердечного ритма является наиболее частым показателем данных изменений. В результате такие пациенты отказываются от визитов к стоматологу, что приводит к ухудшению состояния полости рта. Одной из альтернатив является проведение стоматологических процедур у неадаптированных пациентов под анестезией или седацией. Однако на сегодняшний день отсутствует единый проверенный алгоритм диагностики уровня тревожности у детей и методов ее коррекции для обеспечения безопасного и эффективного лечения.

**ЦЕЛЬ ИССЛЕДОВАНИЯ.** Клиническое и функциональное обоснование выбора метода коррекции тревожности у детей в возрасте 4–12 лет при амбулаторном стоматологическом приеме.

**МАТЕРИАЛЫ И МЕТОДЫ.** В период с 2019 по 2024 г. было проведено исследование с участием 986 детей в возрасте от 4 до 12 лет. До начала обследования осуществлялся сбор общего соматического и психологического анамнеза. Уровень тревожности у детей 4–6 лет определяли с помощью теста детской тревожности Р. Теммля, М. Дорки и В. Амена, а у детей 7–12 лет – с использованием шкалы явной тревожности (CMAS) в адаптации А.М. Прихожан. Гемодинамические параметры (ЧСС и SpO<sub>2</sub>) измеряли детским пульсоксиметром Choice MMed 300C5. Всем пациентам проводился стоматологический осмотр полости рта с определением степени активности кариеса по методике Т.Ф. Виноградовой. На основании полученных данных принималось решение о дальнейшем ведении пациента: проведение лечения при поддержке анестезиолога либо использование арт-терапии в качестве метода психоэмоциональной коррекции.

**РЕЗУЛЬТАТЫ.** По данным психологического тестирования пациенты были распределены на три группы: с низким уровнем тревожности (19,2%), со средним уровнем (52,4%) и с высоким уровнем тревожности (28,4%). У большинства детей с низким уровнем тревожности частота сердечных сокращений находилась в пределах возрастной нормы (86,9%). В этой группе также зафиксирован наибольший процент пациентов с компенсированной активностью кариеса (51,9%). У детей с высоким уровнем тревожности отмечались умеренная и выраженная тахикардия (38,8% и 61,2% соответственно), а также декомпенсированная степень активности кариеса у 62,2% обследованных. Пациенты с низким уровнем тревожности реже нуждались в психологической коррекции по сравнению с детьми со средним и высоким уровнями тревожности. На основании результатов исследования была разработана компьютерная программа «Определение метода коррекции тревожности у детей перед стоматологическим лечением», позволяющая врачам выбирать тактику ведения пациентов перед проведением стоматологических манипуляций.

**ВЫВОДЫ.** Таким образом, для повышения качества и безопасности стоматологической помощи детям на основе комплексного психологического и функционального обследования разработана цифровая программа «Определение метода коррекции тревожности у детей перед стоматологическим лечением».

**Ключевые слова:** тревожность, адаптация, частота сердечных сокращений, арт-терапия, степень активности кариеса.

**Информация о статье:** поступила – 06.03.2025; исправлена – 27.04.2025; принята – 15.05.2025

**Конфликт интересов:** авторы сообщают об отсутствии конфликта интересов.

**Благодарности:** финансирование и индивидуальные благодарности для декларирования отсутствуют.

**Для цитирования:** Никольская И.А., Анисимова Е.Н., Копецкий И.С., Анисимова Н.Ю., Воронин П.А., Катюхина В.А. Выбор метода коррекции тревожности у детей в возрасте 4–12 лет перед стоматологическим лечением. *Эндодонтия Today*. 2025;23(2):312–317. <https://doi.org/10.36377/ET-0089>

## INTRODUCTION

Fear and anxiety before dental treatment are experienced by many people around the world. This is a normal reaction of the body to threatening or unknown influences, which helps the body to mobilize psychological reserves. But for children, going to the dentist's office is very stressful, which they can't cope with on their own. Many studies show that the prevalence of dental anxiety among children varies from 5 to 20% [1–4].

The etiology of anxiety is multi-factorial and consists of personal and indirect experience, psychological characteristics, susceptibility to stressful situations, etc. [5]. The clinical manifestation of anxiety consists of emotional, motor neurohumoral, and vegetative-somatic components [6]. Emotional includes strong subjective experiences, which some children vividly demonstrate by crying and shouting, others by isolation and detachment. Increased muscle tone, motor arousal, or, conversely, blockage of muscle activity are motor manifestations. Hormonal changes in anxiety are characterized by increased levels of ACTH, cortisol, and other hormones [7]. The autonomic-somatic component is expressed in changes in the functioning

of organ systems that are controlled by the sympathetic and parasympathetic nervous systems. An increase in the heart rate is the most common indicator of anxiety and fear [8]. In a study by M.P. Shindova et al., we found a correlation indicating that the degree of self-reported anxiety as a subjective method of assessing psychological state is comparable to the heart rate as an indicator of stress [9]. In the study of Z. Alghareeb et al. significant increases in heart rate were also noted in patients with high dental anxiety [10].

All of the above reactions can be observed even before entering the dentist's office. Patients don't make contact with the doctor, and in the case of compulsory treatment, they form a negative experience, thereby increasing dental anxiety. As a result, children often refuse to attend preventive dental appointments, which leads to poor oral health [11].

Alternatively, these patients are treated under anesthesia or sedation. But, do not forget that the anesthetic risk exceeds the risk from dental intervention. The goal of every dentist should be to treat patients safely and effectively. The problem is that there is no verified algorithm for dentists to determine the level of anxiety of

a child, and in which case it is necessary to prescribe dental treatment under general anesthesia.

It is customary to use psychological testing to diagnose children's anxiety. There are many different questionnaires and scales available in the literature. For a quick and accurate diagnosis, you should choose tests that are suitable for the child's age, simple and understandable in performing and analyzing the result, as well as those that have passed standardization in Russia. Psychological methods should be supported by objective indicators (for example, heart rate data). Only on the basis of a comprehensive diagnosis of the patient will we be able to choose further tactics for working with them.

### PURPOSE OF THE STUDY

Clinical and functional justification of the choice of a method for correcting anxiety in children aged 4–12 years in an outpatient dental appointment

### MATERIALS AND METHODS

From 2019–2024 years were examined 986 children (482 boys and 504 girls) aged 4–12 years, including 472 children aged 4 to 6 years and 514 children aged 7 to 12 years. The inclusion criterion for the study was children aged 4–12 years with varying degrees of anxiety before dental treatment. Non-inclusion criteria include the presence of bronchial asthma, diabetes mellitus, mental retardation, children with physical disabilities and an established psychiatric diagnosis. The exclusion criterion is unwillingness to continue participating in the study.

Before starting the examination, the general somatic and psychological history of patients was collected using developed questionnaires for parents, consisting of key questions about the child's condition and behavior when visiting a dentist.

The next step was to determine the patient's level of anxiety. For children from 4 to 6 years of age, the "Child Anxiety Test" was chosen by R. Temml, M. Dorika, and V. Amen. Children aged from 7 to 12 years were assessed for their level of anxiety using the "Explicit Anxiety Scale CMAS" adapted by A. M. Prikhozhan. According to the results of psychological testing, all patients were divided into three groups: those with a low level of anxiety, medium and high.

The heart rate and blood oxygen saturation level were determined using ChoiceMMeda ChoiceMMed 300C5 pediatric pulse oximeter. The correspondence of the obtained parameters with the age norm, moderate or severe tachycardia was checked according to the protocol of the CSSSA of the FMBA of Russia.

All patients underwent a dental examination of the oral cavity, the CPI index was calculated, and the degree of caries activity was determined by T.F. Vinogradova.

After analyzing the results of all methods of examination, a decision was made on the further algorithm of actions for safe and effective correction of anxiety and rehabilitation of the oral cavity of patients. As a method of psychological correction, was chosen art therapy, using neurographics – a method of organizing thinking


and transforming the psycho-emotional state. Using markers, colored markers, and pencils, the child spent 15–20 minutes drawing neurographic lines and coloring the resulting fragments, thereby forming new neural connections.

The Wilcoxon test was used to determine statistical differences. Differences between independent samples were compared using nonparametric Mann-Whitney criteria and two-sample Kolmogorov-Smirnov criteria.

### RESULTS


According to the results of R. Temml's "Child Anxiety Test" and the "CMAS Scale of explicit anxiety CMAS", a low level of anxiety was detected in 19.2% of the subjects, an average level was observed in 52.4% and a high level in 28.4%, respectively (Fig. 1).

In the group of children with a low level of anxiety, only 13.1% had moderate tachycardia, while the remaining 86.9% had heart rate indicators within the age norm. In this group, the compensated degree of caries activity was found in 51.9%, subcompensated in 33.3%, and decompensated was the lowest percentage 14.8%.




**Fig. 1.** Distribution of children according to the results of psychological testing

**Рис. 1.** Распределение детей в соответствии с результатами психологического тестирования



**Fig. 2.** Comparison of heart rate indicators in children with different levels of anxiety

**Рис. 2.** Сравнение показателей частоты сердечных сокращений у детей с различными уровнями тревожности



**Fig. 3.** Comparison of the degree of caries activity in children with different levels of anxiety

**Рис. 3.** Сравнение степени активности кариеса у детей с различными уровнями тревожности

Among children with an average level of anxiety, the heart rate was within the normal range in 10.1%. 56.8% showed moderate tachycardia, and 33.1% severe. The compensated degree of caries activity was determined in 14.1%, subcompensated in 54.4%, and decompensated in 31.5%.

The group with a high level of anxiety had the highest percentage of patients with severe tachycardia – 61.2%, the remaining – 38.8% had moderate tachycardia. Also in this group, the largest percentage of children with a decompensated degree of caries activity was identified – 62.2%. 28.9% had a subcompensated degree of caries activity and only 8.9% had a compensated one. Fig. 2 and 3 graphically represent the results obtained in groups with low, medium, and high levels of anxiety.

After performing all diagnostic manipulations, analyzing the results and distributing patients into groups, a decision was made on the further tactics of anxiety correction. Children with a high level of anxiety, pronounced tachycardia and decompensated degree of caries activity were referred for further treatment accompanied by an anesthesiologist. The remaining groups of patients underwent correction of their psycho-emotional state by neurography. Children with low levels of an-

xiety needed less adaptation techniques than those with medium and high levels. But, as a result of psychological work, full cooperation of patients and parents with a dentist was achieved, anxiety levels were reduced, and hemodynamic parameters were normalized.

## DISCUSSION

Based on our study, to standardize the protocol for choosing a method for correcting anxiety in children of different age groups, a computerized program “Determining the method for correcting anxiety in children before dental treatment” was developed [12]. The main task of the program is to determine the future strategy of working with the patient based on the collection of anamneses and the results of objective examination methods (how many adaptation techniques should be carried out, what methods should be used, and in what case the patient should be referred for general anesthesia). For each patient in the program, an electronic card is created, which indicates the main contact information about the child and his parents. A questionnaire for collecting general somatic and psychological medical history is built into the program, and parents can fill it out directly in it. The doctor enters the results of determining the level of anxiety, hemodynamic parameters and the CPI index independently in the corresponding columns. After these simple manipulations, the program automatically converts the data according to its internal score system and displays the result on the screen. Also, depending on the answers, when filling out the patient’s general medical history questionnaire, the program can generate a comment about the need to consult with a pediatrician or anesthesiologist to safely choose the method of anesthesia in dental treatment.

## CONCLUSION

Thus, on the basis of a comprehensive psychological and functional examination of children before dental treatment, it became possible to create a digital program: “Determining the method of correcting anxiety in children before dental treatment”, which allows improving the quality and safety of dental care provided to children. The advantage of the developed program is its ease of use, reduced time costs, and increased efficiency in providing dental care to the children’s population.

## REFERENCES / СПИСОК ЛИТЕРАТУРЫ

1. Baier K., Milgrom P., Russell S., Mancil L., Yoshida T. Children’s fear and behavior in private pediatric dentistry practices. *Pediatr Dent*. 2004;26(4):316–321.
2. Alsadat F.A., El-Housseiny A.A., Alamoudi N.M., Elderwi D.A., Ainos A.M., Dardeer F.M. Dental fear in primary school children and its relation to dental caries. *Niger J Clin Pract*. 2018;21(11):1454–1460. [https://doi.org/10.4103/njcp.njcp\\_160\\_18](https://doi.org/10.4103/njcp.njcp_160_18)
3. Zhu M., Yu H., Xie B., Li H., He Q., Li H. Et al. Experiential learning for children’s dental anxiety: a cluster randomized trial. *BMC Oral Health*. 2020;20(1):216. <https://doi.org/10.1186/s12903-020-01204-5>
4. Yon M.J.Y., Chen K.J., Gao S.S., Duangthip D., Lo E.C.M., Chu C.H. Dental fear and anxiety of kindergarten children in Hong Kong: A cross-sectional study. *Int J Environ Res Public Health*. 2020;17(8):2827. <https://doi.org/10.3390/ijerph17082827>
5. Armfield J.M., Heaton L.J. Management of fear and anxiety in the dental clinic: a review. *Aust Dent J*. 2013;58(4):390–407. <https://doi.org/10.1111/adj.12118>
6. Bobrov A.E., Usatenko E.V. The concept of anxiety disorders: main development trends. *Social and Clinical Psychiatry*. 2021;31(4):62–70. (In Russ.).  
Бобров А.Е., Усатенко Е.В. Концепция тревожных расстройств: основные тенденции развития. *Социальная и клиническая психиатрия*. 2021;31(4):62–70.
7. Andrews G., Hobbs M.J., Borkovec T.D., Beesdo K., Craske M.G., Heimberg R.G. et al. Generalized worry

- disorder: a review of DSM-IV generalized anxiety disorder and options for DSM-V. *Depress Anxiety*. 2010;27(2):134–147. <https://doi.org/10.1002/da.20658>
8. Farhat-McHayleh N., Harfouche A., Souaid P. Techniques for managing behaviour in pediatric dentistry: comparative study of live modelling and tell-show-do based on children's heart rates during treatment. *J Can Dent Assoc*. 2009;75(4):283.
  9. Shindova M.P., Belcheva-Krivorova A., Taralov Z. Pulse oximetry in paediatric dentistry. *Folia Med*. 2022;64(2):202–206. <https://doi.org/10.3897/folmed.64.e69136>
  10. Alghareeb Z., Alhaji K., Alhaddad B., Gaffar B. Assessment of dental anxiety and hemodynamic changes during different dental procedures: A report from Eastern Saudi Arabia. *Eur J Dent*. 2022;16(4):833–840. <https://doi.org/10.1055/s-0041-1740222>
  11. Nikolskaia I.A., Katyukhina V.A., Pogabalo I.V., Patrakova N.N., Ogloblin A.A., Kuliev R.M. Basics of formation and correction of dental anxiety. *Endodontics Today*. 2023;21(4):276–280. (In Russ.) <https://doi.org/10.36377/1683-2981-2023-21-4-276-280>
  12. Anisimova E.N., Katyukhina V.A., Nikolskaya I.A., Anisimova N.Yu., Khodnenko O.V. *Determination of the method of anxiety correction in children before dental treatment*. Russian Federation patent No. 2024683372. Publ. 14.10.2024. (In Russ.)
- Анисимова Е.Н., Катюхина В.А., Никольская И.А., Анисимова Н.Ю., Ходненко О.В. *Определение способа коррекции тревожности у детей перед стоматологическим лечением*. Патент Российской Федерации № 2024683372. Оpubл. 14.10.2024.

## INFORMATION ABOUT THE AUTHORS

**Irina A. Nikolskaya** – Cand. Sci. (Med.), Professor, Department of Therapeutic Dentistry, Institute of Dentistry, Pirogov Russian National Research Medical University, 1 Ostrovityanova Str., Moscow 117997, Russian Federation; <https://orcid.org/0000-0001-8042-2884>

**Evgenia N. Anisimova** – Cand. Sci. (Med.), Associate Professor, Professor of the Department of Dentistry and ChLH, Russian Biotechnological University, 11 Volokolamskoe Highway, Moscow 125080, Russian Federation; <https://orcid.org/0000-0001-7109-6431>

**Igor S. Kopetskiy** – Dr. Sci. (Med.), Professor, Head of the Department of Therapeutic Dentistry at the Institute of Dentistry, Pirogov Russian National Research Medical University, 1 Ostrovityanova Str., Moscow 117997, Russian Federation; <https://orcid.org/0000-0002-4723-6067>

**Natalia Y. Anisimova** – Cand. Sci. (Med.), Associate Professor, Associate Professor of the Department of Dentistry and ChLH, Russian Biotechnological University, 11 Volokolamskoe Highway, Moscow 125080, Russian Federation; <https://orcid.org/0000-0002-3073-7041>

**Pavel A. Voronin** – Cand. Sc. (Med.), Associate Professor of the Department of Maxillofacial Surgery and Dentistry of the Institute of Motherhood and Childhood, Pirogov Russian National Research Medical University, 1 Ostrovityanova Str., Moscow 117997, Russian Federation; <https://orcid.org/0000-0002-2134-2320>

**Valeria A. Katyukhina** – Assistant of the Department of Therapeutic Dentistry at the Institute of Dentistry, Pirogov Russian National Research Medical University, 1 Ostrovityanova Str., Moscow 117997, Russian Federation; <https://orcid.org/0009-0004-5092-4997>

## ИНФОРМАЦИЯ ОБ АВТОРАХ

**Никольская Ирина Андреевна** – к.м.н., профессор кафедры терапевтической стоматологии Института стоматологии, ФГАОУ ВО «Российский национальный исследовательский медицинский университет имени Н.И. Пирогова», 117997, Российская Федерация, г. Москва, ул. Островитянова, д. 1; <https://orcid.org/0000-0001-8042-2884>

**Анисимова Евгения Николаевна** – к.м.н., доцент, профессор кафедры стоматологии и челюстно-лицевой хирургии, ФГБОУ ВО «Российский биотехнологический университет (РОСБИОТЕХ)»; 125080, Российская Федерация, г. Москва, Волоколамское шоссе, д. 11; <https://orcid.org/0000-0001-7109-6431>

**Копецкий Игорь Сергеевич** – д.м.н., профессор, заведующий кафедрой терапевтической стоматологии Института стоматологии, ФГАОУ ВО «Российский национальный исследовательский медицинский университет имени Н.И. Пирогова», 117997, Российская Федерация, г. Москва, ул. Островитянова, д. 1; <https://orcid.org/0000-0002-4723-6067>

**Анисимова Наталья Юрьевна** – к.м.н., доцент, доцент кафедры стоматологии и челюстно-лицевой хирургии, ФГБОУ ВО «Российский биотехнологический университет (РОСБИОТЕХ)»; 125080, Российская Федерация, г. Москва, Волоколамское шоссе, д. 11; бывший президент Международной федерации стоматологических анестезиологов (IFDAS); Российская Федерация; <https://orcid.org/0000-0002-3073-7041>

**Воронин Павел Анатольевич** – к.м.н., доцент кафедры челюстно-лицевой хирургии и стоматологии института материнства и детства, ФГАОУ ВО «Российский национальный исследовательский медицинский университет имени Н.И. Пирогова», 117997, Российская Федерация, г. Москва, ул. Островитянова, д. 1; <https://orcid.org/0000-0002-2134-2320>

**Катюхина Валерия Андреевна** – ассистент кафедры терапевтической стоматологии Института стоматологии, ФГАОУ ВО «Российский национальный исследовательский медицинский университет имени Н.И. Пирогова», 117997, Российская Федерация, г. Москва, ул. Островитянова, д. 1; <https://orcid.org/0009-0004-5092-4997>

**AUTHOR'S CONTRIBUTION**

Irina A. Nikolskaya – has made a substantial contribution to the concept or design of the article; revised the article critically for important intellectual content; approved the version to be published.

Evgenia N. Anisimova – has made a substantial contribution to the concept or design of the article; the acquisition, analysis, or interpretation of data for the article; drafted the article; approved the version to be published, drafted the article.

Igor S. Kopetskiy – has made a substantial contribution to the concept or design of the article; revised the article critically for important intellectual content; approved the version to be published.

Natalia Y. Anisimova – has made a substantial contribution to the concept or design of the article; revised the article critically for important intellectual content; approved the version to be published, drafted the article.

Pavel A. Voronin – has made a substantial contribution to the concept or design of the article; revised the article critically for important intellectual content; approved the version to be published, drafted the article.

Valeria A. Katyukhina – the acquisition, analysis, or interpretation of data for the article, drafted the article.

**ВКЛАД АВТОРОВ**

И.А. Никольская – существенный вклад в замысел и дизайн исследования, критический пересмотр статьи в части значимого интеллектуального содержания; окончательное одобрение варианта статьи для опубликования.

Е.Н. Анисимова – существенный вклад в замысел и дизайн исследования, сбор данных, анализ и интерпретация данных, подготовка статьи, окончательное одобрение варианта статьи для опубликования.

И.С. Копецкий – существенный вклад в замысел и дизайн исследования, критический пересмотр статьи в части значимого интеллектуального содержания; окончательное одобрение варианта статьи для опубликования.

Н.Ю. Анисимова – существенный вклад в замысел и дизайн исследования, сбор данных, анализ и интерпретация данных, подготовка статьи, окончательное одобрение варианта статьи для опубликования.

П.А. Воронин – существенный вклад в замысел и дизайн исследования, сбор данных, анализ и интерпретация данных, подготовка статьи, окончательное одобрение варианта статьи для опубликования.

В.А. Катюхина – сбор, анализ или интерпретация данных для статьи, подготовка статьи.