



https://doi.org/10.36377/ET-0116

An ex vivo antimicrobial evaluation after the preparation with XP-Endo Shaper and Trunatomy systems

Alana Cassia Soares Moraes Souza¹ [0], Carlos Eduardo da Silveira Bueno¹ [0], Carlos Eduardo Fontana² [0], Carlos Henrique Meloni¹ [0], Carolina Pessoa Stringheta¹ [0], Alexandre Sigrist De Martin¹ [0], Rina Andrea Pelegrine¹ [0], Wayne Martins Nascimento¹ [0], Ana Grasiela da Silva Limoeiro³ [0] \bowtie , Monique Aparecida de Lima Rios Pitzschk⁴ [0], Aida Meto⁵ [0], Michel Klymus¹ [0], Marilia Fagury Videira Marceliano-Alves^{6,7,8} [0], Daniel Guimarães Pedro Rocha⁵ [0]

□ grasielalimoeiro@gmail.com

Abstract

AIM. The aim of this study was to evaluate bacterial reduction in root canals with the XP-Endo Shaper (XP) and Trunatomy (TN) systems.

MATERIALS AND METHODS. Twenty-eight permanent human type I, oval-shaped Vertucci premolars and straight root canals were contaminated with Enterococcus faecalis for 30 days at 37°C. Samples were collected prior to instrumentation. The teeth were divided into two groups (n = 14) and processed with the tested groups at 37°C: XP group – (30/0.04) and TN group – Small (20/0.04) and Prime (26/0.04). Biological samples before and after instrumentation were collected using a sterile paper cone inserted into the canal for one minute. Bacteria were counted using colony forming units (CFU/mL) and results were subjected to Kruskal-Wallis test at 5 level of significance.

RESULTS. Both the XP and TN systems significantly reduced bacterial counts (p < 0.0001), but did not eliminate bacteria in the root canals.

CONCLUSIONS. Both the Trunatomy and XP-Endo Shaper systems were similar in terms of antimicrobial efficacy, but neither system was able to eliminate bacteria from the root canals.

Keywords: endodontics, Enterococcus faecalis, root canal treatment

Article info: received – 02.06.2025; revised – 03.07.2025; accepted – 24.07.2025

Conflict of interest: The authors report no conflict of interest.

Acknowledgements: This study was supported by grants from Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazilian Governmental Institutions.

For citation: Souza A.C.S.M., Bueno C.E.S., Fontana C.E., Meloni C.H., Stringheta C.P., Martin A.S., Pelegrine R.A., Nascimento W.M., Limoeiro A.G.S., Pitzschk M.A.L.R., Meto A., Klymus M., Marceliano-Alves M.F.V., Rocha D.G.P. An ex vivo antimicrobial evaluation after the preparation with XP-Endo Shaper and Trunatomy systems. *Endodontics Today.* 2025;23(3):441–449. https://doi.org/10.36377/ET-0116





¹ Instituto de Pesquisas São Leopoldo Mandic, Campinas, São Paulo, Brazil

² PUC Campinas, Campinas, São Paulo, Brazil

³ University of Sao Paulo, Bauru, Brazil

⁴ Educational Society University of Santa Catarina, Joinville, Brazil

⁵ School of Dentistry, University of Modena and Reggio Emilia, Italy

⁶ Maurício de Nassau University Centre (UNINASSAU), Rio de Janeiro, Brazil

⁷ Dr. D.Y. Patil Dental College and Hospital, Dr. D.Y. Patil, Pune, India

⁸ Iguaçu University, Nova Iguaçu, Rio de Janeiro, Brazil

⁹ Pontifical Catholic University of Campinas, São Paulo, Brazil

Экспериментальная ex vivo оценка антимикробной эффективности после препарирования с использованием систем XP-Endo Shaper и TruNatomy

А.К.С.М. Соуза¹ (D), К.Э.С. Буэно¹ (D), К.Э. Фонтана² (D), К.Э. Мелони¹ (D), К.П. Стрингета¹ (D), А.С. Мартин¹ (D), Р.А. Пелегрине¹ (D), У.М. Насименто¹ (D), А.Г.С. Лимойру³ (D) ⊠, М.А.Л.Р. Питшк⁴ (D), А. Мето⁵ (D), М. Климус¹ (D), М.Ф.В. Марселиану-Алвес^{6,7,8} (D), Д.Г.П. Роша⁰ (D)

- ¹ Институт исследований São Leopoldo Mandic, Кампинас, Сан-Паулу, Бразилия
- ² Папский католический университет Кампинаса (PUC Campinas), Кампинас, Сан-Паулу, Бразилия
- ³ Университет Сан-Паулу, Бауру, Бразилия
- 4 Университетская образовательная ассоциация Санта-Катарины, Жоинвили, Бразилия
- 5 Стоматологический факультет, Университет Модены и Реджо-Эмилии, Италия
- ⁶ Университетский центр Maurício de Nassau (UNINASSAU), Рио-де-Жанейро, Бразилия
- 7 Стоматологический колледж и госпиталь доктора Д.Я. Патила, Университет доктора Д.Я. Патила, Пуне, Индия
- 8 Университет Игуасу, Нова-Игуасу, Рио-де-Жанейро, Бразилия
- 9 Папский католический университет Кампинаса, Сан-Паулу, Бразилия

☐ grasielalimoeiro@gmail.com

Резюме

ЦЕЛЬ. Целью данного исследования было оценить степень снижения бактериальной обсемененности в корневых каналах после обработки с использованием систем XP-Endo Shaper (XP) и TruNatomy (TN). МАТЕРИАЛЫ И МЕТОДЫ. Двадцать восемь постоянных человеческих премоляров типа I по Вертуччи с овальной формой и прямыми корневыми каналами были заражены $Enterococcus\ faecalis\$ в течение 30 дней при температуре 37°C. Образцы собирались до начала инструментальной обработки. Зубы были разделены на две группы (n=14) и обработаны при 37°C: группа XP – инструмент (30/0.04); группа TN – инструменты Small (20/0.04) и Prime (26/0.04). Биологические образцы до и после обработки собирались стерильным бумажным штифтом, введенным в канал на одну минуту. Подсчет бактерий производился по числу колониеобразующих единиц (КОЕ/мл), результаты анализировались с использованием критерия Крускала—Уоллиса при уровне значимости 5%.

РЕЗУЛЬТАТЫ. Обе системы – XP и TN – достоверно снижали количество бактерий в корневых каналах (p < 0.0001), однако не обеспечивали полного их удаления.

ВЫВОДЫ. Системы TruNatomy и XP-Endo Shaper продемонстрировали сопоставимую антимикробную эффективность, однако ни одна из них не обеспечила полной элиминации бактерий из корневых каналов.

Ключевые слова: эндодонтия, Enterococcus faecalis, лечение корневых каналов

Информация о статье: поступила – 02.06.2025; исправлена – 03.07.2025; принята – 24.07.2025

Конфликт интересов: авторы сообщают об отсутствии конфликта интересов.

Благодарности: Данное исследование было поддержано грантами Фонда имени Карлоса Чагаса Фильо по поддержке научных исследований штата Рио-де-Жанейро (FAPERJ) и Национального совета по-научному и технологическому развитию (CNPq) – государственных научных учреждений Бразилии.

Для цитирования: Соуза А.К.С.М., Буэно К.Э.С., Фонтана К.Э., Мелони К.Э., Стрингета К.П., Мартин А.С., Пелегрине Р.А., Насименто У.М., Лимойру А.Г.С., Питшк М.А.Л.Р., Мето А., Климус М., Марселиану-Алвес М.Ф.В., Роша Д.Г.П. Экспериментальная ех vivo оценка антимикробной эффективности после препарирования с использованием систем XP-Endo Shaper и TruNatomy. *Эндодонтия Today.* 2025;23(3):441–449. https://doi.org/10.36377/ET-0116

INTRODUCTION

One of the main aims of endodontic treatment is to prevent or promote the healing of apical periodontitis by cleaning, shaping, disinfecting and filling the root canal system [1]. The microorganisms can organize themselves into biofilms, which increases their resistance to endodontic procedures and makes root canal disinfection a challenge [2].

One microorganism of interest in endodontics is Enterococcus faecalis, which has been extensively studied as it is particularly associated with persistent endodontic infections, which are a common cause of treatment failure due to their resistance in intracanal procedures [3]. This bacterium has an inherent resistance to various treatments, including irrigation solu-

tions, intracanal medications [4], antibiotics and pH changes [5]. Its ability to adhere, multiply, invade, resist host defenses and compete with other bacteria increases its virulence [6].

Mechanical preparation is the most critical phase of endodontic treatment, and great development has been made in the development of instruments [7]. Studies show that even with different techniques, complete removal of organic debris [8] and bacteria [9] is difficult to achieve. Automated systems based on a nickel-titanium alloy that work with continuous rotary motion have become widely accepted in recent decades [9]. These files, with innovative designs, manufacture, alloy composition and heat treatment [10], have redefined root canal instrumentation.

Some of these systems are XP-Endo Shaper (FKG Dentaire, La Chaux de Fonds, Switzerland) and TruNatomy (Dentsply Sirona, Ballaigues, Switzerland). These files have an innovative design and use rotary motion to enhance the root can preparations [7], considered effective for complex anatomy that often harbor resistant microorganisms such as *Enterococcus faecalis* [5; 6].

The XP-Endo Shaper (FKG) with its serpentine design and NiTi MaxWire alloy adapts to the irregularities of the canal and minimizes stress on the dentin wall. It transitions from a straight, martensitic shape to a spoon shape at body temperature and uses its elasticity and shape memory effect to expand and contract during rotation to improve cleaning and disinfection. This dynamic ability improves root canal treatments by reaching hard-to-reach areas and effectively agitating the irrigant [7; 11]. The instrument is 30/.01 (size/taper), but can have a final canal preparation of 30/.04 [12].

Another new system is the Trunatomy (Dentsply Sirona), which ensures superior peri-cervical dentin preservation thanks to a special instrument geometry and heat treatment [13]. The Trunatomy instruments are characterized by flexibility, regressive tapering and a decentralized parallelogram cross-section and faithfully reproduce the anatomy of the canal [14]. This system comprises five instruments for cervical preparation (Orifice Modifier 20.08), the glide path (Glider 17.02v) and three sizes for canal shaping: Small (20.04v), Prime (26.04v) and Medium (36.03v) [11].

Endo Shaper (FKG Dentaire) and the Trunatomy (Dentsply Sirona) system. The null hypothesis tested was that there would be no difference between the groups in terms of bacterial reduction [15; 16].

MATERIALS AND METHODS

The study was conducted in accordance with the Declaration of Helsinki. This study approved by the local ethics committee because it involved biological material (Opinion: 5.351.873, 14th April, 2022). The manuscript was written according to the Preferred Reporting Items for Laboratory studies in Endodontology (PRILE) 2021 guidelines [17] (Fig. 1).

The number of 14 samples per group was based on previous works [18; 19]. The sample was calculated using the t-test, with a minimum difference between the means of the treatments of 0.74, a standard deviation of 0.81–0.78, a number of treatments of 2, a test power of 0.80 and an alpha of 0.05.

Sample selection and standardization

The criteria included the selection of permanent human mandibular premolars with fully developed, singular roots, a straight single canal curvature < 10° according to Schneider [20] and a Vertucci type I and an oval shape (in which the buccolingual diameter is twice as large as the mesiodistal diameter in the first two thirds of the canal). Teeth with fractures, calcifications, dilacerations or previous endodontic treatment were excluded. Radiographs were taken in both vestibulo-lingual and mesiodistal orientations to confirm compliance with the inclusion criteria.

Studies have highlighted the effectiveness of TruNatomy in shaping and preserving root canal anatomy with limited research on its effects on reducing intracanal bacteria **AIM AND HYPOTHESIS** Aim: To compare the efficacy of mechanical bacterial reduction between the XP-Endo Shaper (FKG Dentaire) and TruNatomy (Dentspty Sirona) systems Hypothesis: The null hypothesis tested was that there would be no differences between groups in terms of bacteria reduction **ETHICAL APPROVAL** This study was submitted to and approved by the local ethics committee (CAAE 5.351.873) because it involved biological material SAMPLES Twenty-eight permanent human premolars with formed single roots and straight canals EXPERIMENTAL AND CONTROL GROUPS, **INCLUDE INDEPENDENT VARIABLES Group XP** (n = 12): The samples in this group were instrumented with the XP-Endo Shaper rotary system **TN group** (n = 12): The samples in this group were instrumented with the TruNatomy rotary system **OUTCOME ASSESSED** Bacterial reduction in root canals METHOD USED TO ASSESS THE OUTCOME Colony forming units (CFU/mL) **RESULTS** There was no difference in the extent of microbial reduction between the TN and XP systems (p = 0.5623) CONCLUSION(S) It was found that the TruNatomy and XP-Endo Shaper instruments were equivalent in microbial reduction of biofilm consisting of Enterococcus faecalis Neither system was able to eliminate bacteria from the root canals **FUNDING DETAILS** None **CONFLICT OF INTEREST** The authors deny any conflict of interests

RATIONALE / JUSTIFICATION

Fig. 1. Preferred Reporting Items for Laboratory studies in Endodontology (PRILE) 2021 guidelines

Рис. 1. Рекомендуемые элементы отчета для лабораторных исследований в эндодонтии (PRILE) – рекомендации 2021 г.



The teeth were ultrasonically cleaned to remove any residual periodontal ligament or calculus. They were stored in 0.1% thymol until the start of the study. In preparation for the experiments, the roots were thoroughly rinsed under running water for one hour to remove all thymol residues. They were then dried with a stream of air and gauze. The samples were then flattened with a diamond disk attached to a ruler and measured with a conventional ruler, ensuring standardization to a length of 15 mm.

The working length (WL) was determined with a K#10 file (K-File, Dentsply Maillefer, Ballaigues, Switzerland) up to the apical foramen, whereby the WL was determined by subtracting 1 mm. The canal was then enlarged with a K#20 file (K-File, Dentsply Maillefer, Ballaigues, Switzerland) to standardize the original canal diameter and make room for bacterial contamination.

Micro-CT Scanning

All teeth were scanned in a micro-CT device (Sky-Scan 1273; Brucker micro-CT, Kontich, Belgium) at 70 kV, 114 mA, 12 µm pixel size, 360° around the vertical axis, with 0.5 rotation step and 2 average frame and with a 1.0 mm thick aluminum filter. After scanning, the N. Recon v.1.6.9.16 software (Bruker micro-CT) was used for the image's reconstruction (ring artifact correction of 5, a beam hardening correction of 50% and a smoothing of 8). The 3-dimensional volume and surface area were measured using CTAn v1.14.4.1 software (Bruker micro-CT).

The CTAn v1.14.4.1 software (Bruker micro-CT) was also used to measure the major diameter of the buccolingual root canal three millimeters from the apex. Root canals that had a buccolingual distance that was at least three times the mesiodistal distance were considered oval [21]. This procedure allowed the samples to be matched and randomly divided in two experimental groups according to Enterococcus faecalis contamination.

Enclosure of the roots in silicone

To prevent the irrigation solution from leaking through the apical foramen during root preparation, the root apices were first covered with utility wax. These roots were then placed in plastic cubes with 4 cm and a diameter of 3 cm, filled with silicone and finally covered with a thin layer of cyanoacrylate around the root to ensure stability. The teeth were then sterilized into an autoclave at 121°C for 15 minutes and subsequently contaminated with *E. faecalis*.

Contamination of the sample

The target microorganism for the infiltration test was E. faecalis (ATCC-29212), which was cultured and stored in BHI liquid medium (Brain Infusion Hearth – BHI – Difco – Detroit, Michigan, USA) with 20% glycerol. To prepare the inoculum, 100 μ L of the E. faecalis broth was transferred to 2 mL of BHI broth and kept in the oven at 37°C for a maximum of 24 hours. After this time, the broth became turbid, which was compared with the Mac Farland 10 scale (1.0x10 CFU/mL). 37 g of BHI was dis-

solved in 1 liter of demineralized water and distributed into smaller containers. After the BHI was prepared, it was autoclaved for 40 minutes at 121 °C, pH 7.4, 0.2 at 25c with typical (g/liter).

Twenty microliters (μ L) of the suspension at the final concentration was added to the root canals using a pipette. A sterile, absorbent cotton was moistened, and placed in four wells of each cell culture plate. The lid of the plate was closed and sealed with adhesive tape and the whole was incubated in an oven at 37° C with 5% CO₂ for 30 days, when the used absorbent cotton was changed in plates wells [4]. During the contamination period, the bacteria viability was checked every three days and BHI was added daily to keep the strains viable to confirm that the contamination was effective. The container was then opened, and the contaminated teeth were used to start the experiment.

Root canal Instrumentation

Once the biofilm had matured, the individual samples were removed prior to instrumentation and randomly allocated into experimental groups (n = 12):

- Group TruNatomy (TN);
- Group XP Endo Shaper (XP).

Samples were collected in a laminar flow chamber using a sterile cone of absorbent paper No. 20 (Endopoints, Manacapuru, Amazonas, Brazil) immersed in the canal for 1 minute. The samples were then transferred to an Eppendorf tube containing 5 mL of sterile 0.85% saline solution and shaken for 30 seconds (Vortex CP 600 Plus, Phoenix, Araraquara, Brazil). Serial dilutions (10⁻¹, 10⁻², 10⁻³ and 10⁻⁴) of this suspension were prepared and transferred to test tubes. Aliquots of 0.1 mL of the suspension and each dilution were seeded in Petri dishes containing BHI agar. The plates were incubated in an oven with 5% CO₂ at 37°C for 24 hours. The number of CFU/mL per culture plate was then counted before the root canals preparations.

After the antimicrobial samples prior to instrumentation, 5 mL of sterile 0.9% saline solution was used as irrigant, and the canals were instrumented. Root canal preparation was performed at 37°C in a heating cabinet (800-Heater; PlasLabs, Lansing, MI) built into the laminar flow. Instrumentation of the root canal was performed as follows:

Group XP (n = 12): Samples in this group were instrumented with the XP-Endo Shaper rotary system. While the canal was filled with 0.9% saline, a K#15 file (K-File, Dentsply Maillefer, Ballaigues, Switzerland) was inserted third by third up to the WL. The file reached the WL by long, even back and forth movements in the longitudinal direction. The movements were wide and uninterrupted so that the conical shape of the canal was defined. Once the WL was reached, 3 incremental movements were done. The teeth were instrumented with the X-Smart Plus motor (Dentsply Maillefer, Ballaigues, Switzerland) at 800 rpm and 1.5 Ncm. At each instrumented root third, the canal was irrigated with 5 mL of sterile 0.9% saline solution (Farmax, Brazil) using a conventional irrigation syringe (Ultradent, USA) and a 25x4 irrigation needle (Injex, São Paulo, Brazil)

and the glide path was performed with a K#20 hand file. A 30/ 04 gutta-percha cone was used to check the final preparation diameter.

TN group (*n* = 12): The specimens in this group were instrumented with the TruNatomy rotary system. While the canal was filled with 0.9% sterile saline, a K#15 file was inserted third by third up to the WL. Initially, the Orifice Modifier 20.08 instrument was used, followed by the 17.02v, 20.04v, 26.04v and 36/.03 instruments. The torque used was 1.5 Ncm and the speed was 500 rpm with continuous rotation using the X-Smart Plus motor. Each time the instrument was used, three to four in and out movements were performed until the WL was reached.

All procedures were performed in a cabinet at 37°C with a heater (800-Heater; PlasLabs, Lansing, USA) to simulate body temperature. In both groups, each root canal was irrigated with 25 mL of saline after 3 preparation cycles (1 cycle corresponds to 3 back and forth movements).

Sample collection after instrumentation

After the final irrigation, the samples of each group were taken again using sterile paper cones, which remained in the canal for 1 minute. After 1 minute, the cones were removed and stored in a plastic container. The dilutions were prepared in test tubes according to the same protocol as the previous collection.

Statistical analysis

The results were analyzed using BioEstat 5.3 and the Shapiro-Wilk normality test was used. The sample showed non-normal behavior, then Kruskal-Wallis parametric test (Student-Newman-Keuls) was chosen, considering a significance level of 5%.

RESULTS

The previous count showed no significant difference between the groups before the preparation (0.5546). The results showed a significant intragroup microbial reduction after root canal instrumentation with the TN (p < 0.0001) and XP (p < 0.0001) systems. Also, no difference was found in an intergroup comparison between the TN and XP systems (p = 0.5623) (Table 1).

DISCUSSION

The primary goal of endodontic treatment involves thorough cleaning, disinfection and proper root canal obturation [22]. However, the complicated anatomy often harbors microorganisms in the bifurcations, isthmi and dentinal tubules, making them resistant to conventional mechanical and chemical treatments. This resistance could contribute to the failure of endodontic therapy [23].

In the present study, the instrumentation capacity of 2 rotary systems for the removal of E. faecalis was compared. Despite the different designs, cross-sections and heat treatments, there was no difference in the ability to disinfect root canals with the XP-Endo Shaper and the TruNatomy system. Both systems promoted partial bacterial reduction. The null hypothesis was therefore accepted.

Table 1. Kruskal-Wallis (Student-Newman-Keuls) statistical test of colony forming units/mL (log10) of the sample groups

Таблица 1. Результаты статистического анализа методом Крускала-Уоллиса (и критерия Стьюдента – Ньюмена – Келса) количества колониеобразующих единиц/мл (log10) в исследуемых группах

	TN		ХР	
	PC	CAI	PC	CAI
MN	4.28	3.33	5.46	3.38
MX	6.30	4.98	7.05	5.40
MD (ID)	5.98(0.50) ^A	3.97(0.55) ^B	6.00(0.74) ^A	4.09(1.24) ^B
MA (SD)	5.79 (0.52)	4.00 (0.45)	6.08 (0.47)	4.29 (0.68)

Note: TN – TruNatomy system; XP – XP-Endo Shaper system; PC – prior collection; CAI – collection after instrumentation; MN – minimum values; MX – maximum values; MD – median; ID – interquartile deviations; MA – arithmetic means; SD – standard deviation; ρ < 0.0001

Примечание: TN – система TruNatomy; XP – система XP-Endo Shaper; PC – забор до инструментальной обработки; CAI – забор после инструментальной обработки; MN – минимальные значения; MX – максимальные значения; MD – медиана, ID –межквартильные размахи; средние MA – арифметические значения; SD – стандартное отклонение; p < 0.0001

Oval mandibular premolars were selected for this study because cleaning and shaping these teeth is challenging [12]. This is because the canal usually has a round cross-section during rotary instrumentation and the polar areas in oval canals remain unprepared.

In this study, the final preparation size was 30/0.04 for XP-Shaper and 36/0.03 for Trunatomy. Despite the differences in final apical diameter, studies have shown that both systems have similar capabilities in reducing microbial presence in root canals. This consistency of results suggests that factors such as instrument flexibility, cutting efficiency and ability to move irrigants may play a more critical role than preparation size alone [11]. The design and material composition of the instruments facilitate thorough cleaning of the canal so that even complicated canal anatomies can be reached effectively. These results emphasize the sophistication of modern endodontic instruments, where design complements size to achieve the desired results in bacterial reduction [11].

Enterococcus faecalis was focused on in this study because it appears to be the most common microorganism in persistent endodontic infections [24]. This bacterium tolerates hostile alkaline pH values and resists prolonged nutrient deprivation with reduced metabolic activity in treated root canals [25]. In addition, mature E. faecalis biofilms in root canal dentin exhibit greater resistance to disinfectant solutions than young biofilms [26].

In this study, a culture-based microbiological test based on an ex vivo approach was chosen. Despite newer, more advanced methods for evaluating cell cultures and root canal disinfection, this method is widely supported in the literature [11; 27]. Collection via sterile paper tips, followed by serial dilution and seeding in culture media, facilitated the enumeration of colonies in Petri dishes [28]. Although these systems have similarities in potential bacterial reduction in root canals, it is important to emphasize the method chosen for this study, the bacterial counting [27; 28], which is crucial for the evaluation of instruments and substances. which is crucial for the evaluation of instruments and substances. However, it must be acknowledged that the use of paper points may underestimate the bacterial count in the canal system, as bacteria located in areas further away from the main canal lumen, such as dentinal tubules, isthmuses and bifurcations, may not be detected [1].

Other studies investigating bacterial reduction are in agreement with these results [2; 29]. It has been shown that the reciprocating, manual and rotary technique in combination with 2.5% NaOCI is equally effective in reducing the number of microorganisms in the oval root canals [2]. The efficacy of Trunatomy and Rotate in the disinfection of root canals [29]. Although without significant differences, the preparation with XP-Endo Shaper resulted in a significant reduction in

bacteria compared to TruNatomy instruments [14] or compared to compared to Reciproc [30], which contradicts the present study.

The results of the study emphasize the continuing need for research and innovation in endodontics and highlight the importance of developing new instruments, techniques and strategies for thorough root canal disinfection, especially in cases with resistant microorganisms. Although the study provides valuable insights, the limitations of the study caution against immediate application of the results in clinical practice. It suggests that further research, including clinical trials, is needed to validate and build on the observations.

CONCLUSION

It was found that the Trunatomy and XP-Endo Shaper instruments were equivalent in microbial reduction of biofilm consisting of Enterococcus faecalis. Neither system was able to remove bacteria from the root canals. Therefore, the integration of effective irrigation protocols and solutions is crucial to improve the antibacterial efficacy of these systems. Continued research and development are essential to refine these technologies and improve the clinical outcomes of endodontic therapy.

REFERENCES / СПИСОК ЛИТЕРАТУРЫ

- Siqueira J.F. Jr, Rôças I.N. A critical analysis of research methods and experimental models to study the root canal microbiome. *Int Endod J.* 2022;55(Suppl. 1):46–71. https://doi.org/10.1111/iej.13656
- Nakamura V.C., Candeiro G.T., Cai S., Gavini G. Ex vivo evaluation of three instrumentation techniques on E. faecalis biofilm within oval shaped root canals. *Braz Oral Res.* 2015;29(1):1–7. https://doi.org/10.1590/1807-3107BOR-2015.vol29.0027
- Gomes B.P., Berber V.B., Kokaras A.S., Chen T., Paster B.J. Microbiomes of endodontic-periodontal lesions before and after chemomechanical preparation. *J Endod.* 2015;41(12):1975–1984. https://doi.org/10.1016/j.joen.2015.08.022
- Matos Neto M., Santos S.S., Leão M.V., Habitante S.M., Rodrigues J.R., Jorge A.O. Effectiveness of three instrumentation systems to remove Enterococcus faecalis from root canals. *Int Endod J.* 2012;45(5):435–438. https://doi.org/10.1111/j.1365-2591.2011.01994.x
- Evans M., Davies J.K., Sundqvist G., Figdor D. Mechanisms involved in the resistance of Enterococcus faecalis to calcium hydroxide. *Int Endod J.* 2002;35(3):221–228. https://doi.org/10.1046/j.1365-2591.2002.00504.x
- Kayaoglu G., Ørstavik D. Virulence factors of Enterococcus faecalis: relationship to endodontic disease. Crit Rev Oral Biol Med. 2004;15(5):308–320. https://doi. org/10.1177/154411130401500506
- Marceliano-Alves M.F.V., Ronquete V., Coutinho T.M.C., Boukpessi T., Salvioni A.L.F., Goulart P.A.S.R. et al. Unprepared areas and centralization of oval canals prepared with WaveOne Gold or XP-endo Shaper: microcomputed tomographic analyses. *Acta Odontol Latinoam*. 2023;36(3):177–182. https://doi.org/10.54589/aol.36/3/177

- Versiani M.A., Pécora J.D., de Sousa-Neto M.D. Flatoval root canal preparation with self-adjusting file instrument: a micro-computed tomography study. J Endod. 2011;37(7):1002–1007. https://doi.org/10.1016/j.joen.2011.03.017
- Neves M.A.S., Provenzano J.C., Rôças I.N., Siqueira J.F. Jr. Clinical antibacterial effectiveness of root canal preparation with reciprocating single-instrument or continuously rotating multi-instrument systems. J Endod. 2016;42(1):25–29. https://doi.org/10.1016/j.joen.2015.09.019
- Aksoy Ç., Keriş E.Y., Yaman S.D., Ocak M., Geneci F., Çelik H.H. Evaluation of XP-endo Shaper, Reciproc Blue, and ProTaper Universal NiTi systems on dentinal microcrack formation using micro-computed tomography. *J Endod.* 2019;45(3):338–342. https://doi.org/10.1016/j.joen.2018.12.005
- Macedo F.P.G., Soares A.J., Marceliano-Alves M.F.V., Martinez E., Lopes R., Bastos L.F. et al. The effect of root canal preparation tapers on planktonic bacteria and biofilm reduction in the apical third: A correlative microtomography and microbiological laboratory study. *Int Endod J.* 2024;57(6):700–712. https://doi.org/10.1111/ iej.14052
- Lacerda M.F.L.S., Marceliano-Alves M.F., Pérez A.R., Provenzano J.C., Neves M.A.S., Pires F.R. et al. Cleaning and shaping oval canals with 3 instrumentation systems: a correlative micro-computed tomographic and histologic study. *J Endod*. 2017;43(11):1878–1884. https:// doi.org/10.1016/j.joen.2017.06.032
- 13. Elnaghy A.M., Elsaka S.E., Mandorah A.O. In vitro comparison of cyclic fatigue resistance of TruNatomy in single and double curvature canals compared with different nickel-titanium rotary instruments. BMC Oral

- Health. 2020;20:38. https://doi.org/10.1186/s12903-020-1027-7
- 14. Loyola-Fonseca S.C., Campello A.F., Rodrigues R.C.V., Alves F.R.F., Brasil S.C., Vilela C.L.S. et al. Disinfection and shaping of Vertucci class ii root canals after preparation with two instrument systems and supplementary ultrasonic activation of sodium hypochlorite. *J Endod.* 2023;49(9):1183–1190. https://doi.org/10.1016/j. joen.2023.06.017
- 15. Silva E.J.N.L., Lima C.O., Barbosa A.F.A., Lopes R.T., Sassone L.M., Versiani M.A. The impact of Tru-Natomy and ProTaper gold instruments on the preservation of the periradicular dentin and on the enlargement of the apical canal of mandibular molars. *J Endod.* 2022;48(5):650–658. https://doi.org/10.1016/j.joen.2022.02.003
- Oliveira S.C.A., Bueno C.E.D.S., Pelegrine R.A., Fontana C.E., Martin A.S., Stringheta C.P. Debridement ability of the WaveOne Gold and TruNatomy systems in the apical third of root canals: ex vivo assessment. *Braz Dent J.* 2024;35:5773. https://doi.org/10.1590/0103-6440202405773
- Nagendrababu V., Murray P.E., Ordinola-Zapata R., Peters O.A., Rôças I.N., Siqueira J.F. Jr et al. PRILE 2021 guidelines for reporting laboratory studies in Endodontology: A consensus-based development. *Int Endod J.* 2021;54(9):1482–1490. https://doi. org/10.1111/iej.13542
- Colak M., Evcil S., Bayindir Y.Z., Yigit N. The effectiveness of three instrumentation techniques on the elimination of Enterococcus faecalis from a root canal: an in vitro study. *J Contemp Dent Pract*. 2005;6(1):94–106. https:// doi.org/10.5005/jcdp-6-1-94
- Machado M.E., Sapia L.A., Cai S., Martins G.H., Nabeshima C.K. Comparison of two rotary systems in root canal preparation regarding disinfection. *J Endod.* 2010;36(7):1238–1240. https://doi.org/10.1016/j. joen.2010.03.012
- Schneider S.W. A comparison of canal preparations in straight and curved root canals. *Oral Surg Oral Med Oral Pathol*. 1971;32(2):271–275. https://doi. org/10.1016/0030-4220(71)90230-1
- Jou Y.T., Karabucak B., Levin J., Liu D. Endodontic working width: current concepts and techniques. *Dent Clin North Am.* 2004;48(1):323–335. https://doi. org/10.1016/j.cden.2003.12.006

- 22. Haapasalo M., Shen Y., Qian W., Gao Y. Irrigation in endodontics. *Dent Clin North Am.* 2010;54(2):291–312. https://doi.org/10.1016/j.cden.2009.12.001
- 23. Xu J., Gao Y., Meng Y., Wu W., Tsauo C., Guo T. et al. Mechano-chemical coupling of irrigation enhances endodontic biofilm debridement. *Biofouling*. 2020;36(7):792–799. https://doi.org/10.1080/08927014.2020.1814753
- 24. Pinheiro E.T., Gomes B.P., Ferraz C.C., Sousa E.L., Teixeira F.B., Souza-Filho F.J. Microorganisms from canals of root-filled teeth with periapical lesions. *Int Endod J.* 2003;36(1):1–11. https://doi.org/10.1046/j.1365-2591.2003.00603.x
- 25. Stuart C.H., Schwartz S.A., Beeson T.J., Owatz C.B. Enterococcus faecalis: its role in root canal treatment failure and current concepts in retreatment. *J Endod.* 2006;32(2):93–98. https://doi.org/10.1016/j.joen.2005.10.049
- 26.Wang Z., Shen Y., Haapasalo M. Effectiveness of endodontic disinfecting solutions against young and old Enterococcus faecalis biofilms in dentin canals. *J Endod*. 2012;38(10):1376–1379. https://doi.org/10.1016/j.joen.2012.06.035
- 27. Fernandes K.G.C., Silva B.B.D., Boer N.C., Mandarini D.R., Moreti L.C.T., Kato A.S. The effectiveness of three irrigation systems in the enterococcus faecalis reduction after instrumentation with a reciprocating instrument. *Eur J Dent.* 2020;14(4):539–543. https://doi.org/10.1055/s-0040-1714760
- 28. Soares J.A., Santos Soares S.M.C., Santos César C.A., de Carvalho M.A.R., Brito-Júnior M., de Sousa G.R. et al. Monitoring the effectiveness of photodynamic therapy with periodic renewal of the photosensitizer on intracanal Enterococcus faecalis biofilms. *Photodiagnosis Photodyn Ther.* 2016;13:123–127. https://doi.org/10.1016/j.pdpdt.2016.01.002
- 29. Usta S.N., Solana C., Ruiz-Linares M., Baca P., Ferrer-Luque C.M., Cabeo M., Arias-Moliz M.T. Effectiveness of conservative instrumentation in root canal disinfection. *Clin Oral Investig.* 2023;27(6):3181–3188. https://doi.org/10.1007/s00784-023-04929-z
- Alves F.R.F., Paiva P.L., Marceliano-Alves M.F., Cabreira L.J., Lima K.C., Siqueira J.F. Jr et al. Bacteria and hard tissue debris extrusion and intracanal bacterial reduction promoted by XP-endo shaper and reciproc instruments. *J Endod.* 2018;44(7):1173–1178. https://doi.org/10.1016/i.joen.2018.04.007

INFORMATION ABOUT THE AUTHORS

Alana Cassia Soares Moraes Souza – Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, SP, Brazil; https://orcid.org/0009-0000-5389-5264

Carlos Eduardo da Silveira Bueno – Dentist, MSc, PhD and Professor, Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, São Paulo, Brazil; https://orcid.org/0000-0002-2675-0884

Carlos Eduardo Fontana – Dentist, Professor, Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, São Paulo, Brazil; https://orcid.org/0000-0002-2868-6839

Carlos Henrique Meloni – Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, SP, Brazil; https://orcid.org/0009-0004-0642-9067

Carolina Pessoa Stringheta – Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, SP, Brazil; https://orcid.org/0000-0002-0022-558X

Alexandre Sigrist De Martin – Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, SP, Brazil; https://orcid.org/0000-0002-3320-9172

Rina Andrea Pelegrine – Dentist, MSc, PhD and Professor in the Department of Endodontics, Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, São Paulo, Brazil; https://orcid.org/0000-0003-4175-2121



Wayne Martins Nascimento – Dentist, MSc, Phd and Professor and Researcher Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, São Paulo, Brazil; https://orcid.org/0000-0003-4201-4710

Ana Grasiela da Silva Limoeiro – Dentist, MSc, PhD and Professor, Department of Dentistry, Endodontics and Dental Materials, Bauru Dental School, University of Sao Paulo, Bauru, Brazil; https://orcid.org/0000-0003-4633-720X

Monique Aparecida de Lima Rios Pitzschk – Educational Society University of Santa Catarina, Joinville, Brazil; https://orcid.org/0009-0002-2107-5397

Aida Meto – School of Dentistry, University of Modena and Reggio Emilia, Italy; https://orcid.org/0000-0002-3354-2194 **Michel Klymus** – Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Department of Endodontics, Campinas, SP, Brazil; https://orcid.org/0000-0001-6429-7964

Marilia Fagury Videira Marceliano-Alves – Dentist, Holds MSc and PhD degrees in Endodontics, Professor and Researcher, Professor at Posrgraduate Program in Dentistry, Iguaçu University, Nova Iguaçu, Brazil; https://orcid.org/0000-0002-2917-5934

Daniel Guimarães Pedro Rocha – Dr. Sci. (Med.), Lecturer and Researcher in the Department of Endodontics at the Faculty of Dentistry, PUC Campinas, Department of Endodontics, Center of Life Sciences, Programa de pós-graduação em Ciências da Saúde, Campinas, São Paulo, Brazil; https://orcid.org/0000-0001-9792-2260

ИНФОРМАЦИЯ ОБ АВТОРАХ

Алана Кассия Соарес Мораэс Соуза – факультет Сан-Леопольду Мандик, Институт исследований Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0009-0000-5389-5264

Карлос Эдуардо да Силвейра Буэно – врач-стоматолог, профессор, факультет Сан-Леопольду Мандик, Институт исследований Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0000-0002-2675-0884

Карлос Эдуардо Фонтана – стоматолог, профессор, факультет Сан-Леопольду Мандик, Научно-исследовательский институт Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, штат Сан-Паулу, Бразилия; https://orcid.org/0000-0002-2868-6839

Карлос Энрике Мелони – факультет Сан-Леопольду Мандик, Институт исследований Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0009-0004-0642-9067

Каролина Пессоа Стрингета – факультет Сан-Леопольду Мандик, Институт исследований Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0000-0002-0022-558X

Александр Сигрист Де Мартин – факультет Сан-Леопольду Мандик, Институт исследований Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0000-0002-3320-9172

Рина Андреа Пелегрине – д.м.н., преподаватель кафедры эндодонтии, факультет Сан-Леопольду Мандик, Институт исследований Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0000-0003-4175-2121

Уэйн Мартинс Насименто – врач-стоматолог, преподаватель и исследователь, факультет Сан-Леопольду Мандик, Институт исследований Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0000-0003-4201-4710

Ана Гразиэла да Силва Лимойру – врач-стоматолог, магистр в области эндодонтии, кафедра стоматологии, эндодонтии и стоматологических материалов, Стоматологическая школа в Бауру, Университет Сан-Паулу, Бауру, Бразилия; https://orcid.org/0000-0003-4633-720X

Моник Апаресида де Лима Риос Питшк – Университет Санта-Катарины, Жоинвиль, Бразилия; Факультет Сан-Леопольду Мандик, Институт исследований Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0009-0002-2107-5397

Аида Мето – Стоматологическая школа, Университет Модены и Реджо-Эмилии, Италия; https://orcid.org/0000-0002-3354-2194

Мишель Климус – факультет Сан-Леопольду Мандик, Научно-исследовательский институт Сан-Леопольду Мандик, кафедра эндодонтии, Кампинас, штат Сан-Паулу, Бразилия; https://orcid.org/0000-0001-6429-7964

Марилия Фагури Видейра Марселиану-Алвес – врач-стоматолог, имеет степени магистра и доктора философии (MSc и PhD) в области эндодонтии, преподаватель и исследователь, профессор программы последипломного образования по стоматологии, Университет Игуасу (Iguaçu University), Нова-Игуасу, Бразилия; https://orcid.org/0000-0002-2917-5934

Даниэль Гимарайнс Педру Роша – д.м.н., преподаватель и исследователь кафедры эндодонтии стоматологического факультета Университета PUC Campinas, кафедра эндодонтии, Центр наук о жизни, программа последипломного образования по наукам о здоровье, Кампинас, Сан-Паулу, Бразилия; https://orcid.org/0000-0001-9792-2260

AUTHOR'S CONTRIBUTION

Alana Cassia Soares Moraes Souza - data acquisition and analysis, manuscript preparation, manuscript review.

Carlos Eduardo da Silveira Bueno – data analysis, design, definition of intellectual content, manuscript preparation, manuscript review.

 $Carlos\ Eduardo\ Fontana-data\ analysis,\ manuscript\ preparation,\ manuscript\ review.$



Carlos Henrique Meloni – design, definition of intellectual content, manuscript preparation, manuscript review.

Carolina Pessoa Stringheta - design, definition of intellectual content, manuscript preparation, manuscript review.

Alexandre Sigrist De Martin – data analysis, design, definition of intellectual content, manuscript preparation, manuscript review.

Rina Andrea Pelegrine - manuscript preparation, manuscript review.

Wayne Martins Nascimento - data acquisition and analysis, manuscript review.

Ana Grasiela da Silva Limoeiro - data acquisition and analysis, manuscript review.

Monique Aparecida de Lima Rios Pitzschk - data analysis, design, definition of intellectual content, manuscript review.

Aida Meto – design, definition of intellectual content, manuscript review.

Michel Klymus - design, definition of intellectual content, manuscript review.

Marilia Fagury Videira Marceliano-Alves – design, definition of intellectual content, manuscript preparation, manuscript review.

Daniel Guimarães Pedro Rocha - design, definition of intellectual content, manuscript review.

ВКЛАД АВТОРОВ

А.К.С.М. Соуза - сбор и анализ данных, подготовка рукописи, рецензирование рукописи.

К.Э.С. Буэно – анализ данных, дизайн исследования, формирование интеллектуального содержания, подготовка рукописи, рецензирование рукописи.

К.Э. Фонтана – анализ данных, подготовка рукописи, рецензирование рукописи.

К.Э. Мелони – дизайн исследования, формирование интеллектуального содержания, подготовка рукописи, рецензирование рукописи.

К.П. Стрингета – дизайн исследования, формирование интеллектуального содержания, подготовка рукописи, рецензирование рукописи.

А.С. Мартин – анализ данных, дизайн исследования, формирование интеллектуального содержания, подготовка рукописи, рецензирование рукописи.

Р.А. Пелегрине – подготовка рукописи, рецензирование рукописи.

У.М. Насименто – сбор и анализ данных, рецензирование рукописи.

А.Г.С. Лимойру – сбор и анализ данных, рецензирование рукописи.

М.А.Л.Р. Питшк – анализ данных, дизайн исследования, формирование интеллектуального содержания, рецензирование рукописи.

А. Мето – дизайн исследования, формирование интеллектуального содержания, рецензирование рукописи.

Мишель Климус – дизайн исследования, формирование интеллектуального содержания, рецензирование рукописи

М.Ф.В. Марселиану-Алвес – дизайн исследования, формирование интеллектуального содержания, подготовка рукописи, рецензирование рукописи.

Д.Г.П. Роша – дизайн исследования, формирование интеллектуального содержания, рецензирование рукописи.